

**AIRBORNE URANIUM MONITORING
U. S. ARMY POHAKULOA TRAINING AREA
ISLAND OF HAWAII**

Summary Report - May 2009

**Contract No. W9128A-04-D-0019
Task Order 0040**

Prepared for

**U. S. Army Corps of Engineers
Honolulu District
Fort Shafter, Hawaii**

and

**U. S. Army Garrison, Hawaii
Schofield Barracks, Hawaii**

Prepared by:

**J. W. Morrow, DrPH
Environmental Management Consultant
Honolulu, Hawaii**

ACRONYMS AND ABBREVIATIONS

ATSDR	Agency for Toxic Substances and Disease Registry
EPA	U. S. Environmental Protection Agency
ICP-MS	inductively coupled mass spectrometry
IDL	instrument detection limit
lpm	liters per minute
µg	micrograms
µg/m ³	micrograms per cubic meter
MRL	minimal risk level
N	number of samples
ppm	parts per million
PRL	practical reporting limit
PTA	Pohakuloa Training Area
TSP	total suspended particulate matter
U	uranium
²³⁸ U	uranium-238 isotope
²³⁴ U	uranium-234 isotope
²³⁵ U	uranium-235 isotope
WHO	World Health Organization


INTRODUCTION

An airborne uranium monitoring project at the U. S. Army's Pohakuloa Training Area (PTA) commenced on 4 February 2009. Portable samplers operating at a nominal 5 liters per minute (lpm) are located at three (3) sites on PTA (Figure 1). The samplers were originally set to collect total suspended particulate matter (TSP) from midnight to midnight on sample days. However, due to the very low uranium content of the TSP samples, the run time was increased to 72 hours on 19 Apr 09 and has continued at that rate in an effort to raise the collected uranium mass above the practical reporting level (PRL). Since there was no heavy weapons firing activity on PTA ranges in May, EPA's published once-every-six-days schedule was followed with no additional sampling days.

The 47-mm Teflon filters with the collected TSP are sent to laboratories for gravimetric and then uranium analysis. The analysis method for uranium is inductively coupled plasma - mass spectrometry (ICP-MS), a method capable of detecting uranium down to the picogram (10^{-12} gram) level.

Seventeen (17) samples were collected and analyzed during May 2009, and the results are presented herein.

FIGURE 1
MONITORING SITES



RESULTS AND DISCUSSION

The analysis results for each of the three (3) monitoring stations are summarized in Tables 1 and Figures 2 - 4. The figures also indicate the World Health Organization (WHO) and U. S. Agency for Toxic Substances and Disease Registry (ATSDR) guidelines for uranium exposure protection. The WHO guideline is an annual average while the ATSDR guideline is based on chronic exposure (365 days or longer) to highly soluble uranium compounds. It is clear that the uranium concentrations found at PTA in May 2009 are well below both those health guidelines.

TABLE 1
TSP & AIRBORNE URANIUM CONCENTRATIONS
MAY 2009

Station No.	N	TSP Range ($\mu\text{g}/\text{m}^3$)	U Range ($\mu\text{g}/\text{m}^3$)	U Mean ($\mu\text{g}/\text{m}^3$)	ACTIVITY
1	5	6.9 - 29.7	0.000005 - 0.000009	0.000007	No heavy weapons activity
4	6	5.4 - 29.2	0.000003 - 0.000008	0.000005	No heavy weapons activity
14	6	5.2 - 32.7	0.000003 - 0.000007	0.000005	No heavy weapons activity

As noted in previous reports, the total uranium mass found on each filter was well above, e.g., in this data set 6 to 19 times, the laboratory's instrument detection level (IDL) for the ICP-MS method; however, despite the increased sampling time (72 hours), collected uranium mass remained below the practical reporting level (PRL).

The uranium mass in the 72-hour samples did reach as high as 84% of the PRL in one sample and the average of all samples was 48% of the PRL, a 2% increase over the April 09 data. The fact that total uranium levels remain below the PRL despite the tripling of sampling time is likely due to the very low concentrations being measured and the uncertainty associated with the analytical method at such low levels.

The fact that the measured uranium values continue to be less than the PRL remains significant from a public health perspective. At a nominal sampler flow rate of 5 lpm, the laboratory's PRL of 0.00025 microgram (μg) corresponds to an airborne uranium concentration of 0.000035 $\mu\text{g}/\text{m}^3$, a value several orders of magnitude below health effects guidelines. Uranium isotopes ^{234}U and ^{235}U were again undetectable.

FIGURE 2


24-HOUR URANIUM CONCENTRATIONS
STATION 1

FIGURE 3


AIRBORNE URANIUM CONCENTRATIONS
STATION 4

FIGURE 4

AIRBORNE URANIUM CONCENTRATIONS
STATION 14